

SUPPLEMENT

Table S1. *Escherichia coli* strains and plasmids used in the work

Strain or plasmid	Genotype or characteristic	Reference or source
Strains		
W3110	$F^- \lambda^- rpoS(Am) rph-1 Inv(rrnD-rrnE)$	[1]
DY330	W3110 $\Delta lacU169 gal490 pgl\Delta 8 \lambda cI857 \Delta(cro-bioA)$	[2]
XL1-Blue	<i>recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac</i> [F' <i>proAB lacIqZ</i> Δ M15 Tn10 (Tet ^R)]	Stratagene
KNB800	W3110 $\Delta rply::cat$	[3]
MS15	DY330 <i>rply-1::bla</i> (L25 wild-type)	this work
MS20	DY330 <i>rply-2::bla</i> (L25 Y31A)	this work
MS21	DY330 <i>rply-3::bla</i> (L25 H88F)	this work
MD6	DY330 <i>rply-4::bla</i> (L25 S17L/Y31L)	this work
MD7	DY330 <i>rply-5::bla</i> (L25 S17L/H88F)	this work
MD8	DY330 <i>rply-6::bla</i> (L25 S17L/I29F/D90Y)	this work
MD9	DY330 <i>rply-7::bla</i> (L25 S17L/Y31L/H88F)	this work
MS23	W3110 <i>rply-1::bla</i>	this work
MS25	W3110 <i>rply-2::bla</i>	this work
MS26	W3110 <i>rply-3::bla</i>	this work
MD10	W3110 <i>rply-4::bla</i>	this work
MD11	W3110 <i>rply-5::bla</i>	this work
MD12	W3110 <i>rply-6::bla</i>	this work
MD13	W3110 <i>rply-7::bla</i>	this work
Plasmids		
blank pCR-blunt	cloning vector Km ^R Zeo ^R pUCori <i>Plac lacZα</i>	[3]
PKAB101	pCR-blunt/L25 wild-type	[3]
PKAB119	pCR-blunt/L25 Y31A	[4]
PKAB129	pCR-blunt/L25 H88F	[4]
pRP6	pCR-blunt/L25 S17L	this work
pRP7	pCR-blunt/L25 S17L/I29F	this work
pRP8	pCR-blunt/L25 S17L/Y31L	this work
pRP9	pCR-blunt/L25 S17L/H88F	this work
pRP10	pCR-blunt/L25 S17L/I29F/D90Y	this work
pRP11	pCR-blunt/L25 S17L/Y31L/H88F	this work
pBR322	cloning vector Ap ^R Tc ^R	[5]
pUC18	cloning vector Ap ^R <i>Plac lacZα</i> pMB1 ori	US Biological
pUC18/Eco5Sfr	fragment of <i>E. coli</i> 5S rRNA gene cloned into pUC18	this work

Table S2. Oligonucleotides used in this work

Name	Nucleotide sequence (5' → 3')
L25-S17L-F	GTAAGGGTGC GTTACGCCGCCTGC
L25-S17L-R	GCAGGCGGCGTAACGCACCCTTAC
L25-I29F-F	CAAGTTCCCGGCATTTATCTACGGTGGC
L25-I29F-R	GCCACCGTAGATAAATGCCGGGAACTTG
L25-D90Y-F	GCAGCACATCTATTTTCGTTTCGCG
L25-D90Y-R	CGCGAACGAAATAGATGTGCTGC
L25-Y31L-F	GGCAATCATCCTGGGTGGCAAAG
L25-Y31L-R	CTTTGCCACCCAGGATGATTGCC
L25-H88F-F	CCGAAGCTGCAGTTCATCGACTTC
L25-H88F-R	GAAGTCGATGAACTGGAGCTTCGG
AK22	GGTCGCCTGTAGCAGGATTAATTTAAGAGAGAAAGAAATGTTTACTAT CAACGCAGAAG
AK23	AAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACTTAAGCGCGAAC GAAGTCGATG
rplY-D90Y- rec-R	AAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACTTAAGCGCGAAC GAAATAGATGTGC
AK21	CCCGCCGGAGCGGGGTTTTTTACA ACTTATTCAGCAATTACCAATGCTT AATCAGTGAGGCACC
rplY-check-F	GCAGATCACTTACCTGTTGC
rplY-check- R	GTTTTTTACA ACTTATTCAGC
Fr5S-F	GGGAATTCCATATGTAATACGACTCACTATAGCCGATGGTAGTGTGGG
Fr5S-R	CTAGTCTAGACCTACTCTCGCATGGGGAGACC
5S-biotin*	GCGCAGCGAG

* 3'-biotinylated RNA oligonucleotide.

REFERENCES

1. Bachmann, B. J. (1972) Pedigrees of some mutant strains of *Escherichia coli* K-12, *Bacteriol. Rev.*, **36**, 525-557.
2. Yu, D., Ellis, H. M., Lee, E. C., Jenkins, N. A., Copeland, N. G., and Court, D. L. (2000) An efficient recombination system for chromosome engineering in *Escherichia coli*, *Proc. Natl. Acad. Sci. USA*, **97**, 5978-5983.
3. Korepanov, A. P., Gongadze, G. M., Garber, M. B., Court, D. L., and Bubunenko, M. G. (2007) Importance of the 5S rRNA-binding ribosomal proteins for cell viability and translation in *Escherichia coli*, *J. Mol. Biol.*, **366**, 1199-1208.
4. Korobeinikova, A. V., Gongadze, G. M., Korepanov, A. P., Eliseev, B. D., Bazhenova, M. V., and Garber, M. B. (2008) 5S rRNA-recognition module of CTC family proteins and its evolution, *Biochemistry (Moscow)*, **73**, 156-163.
5. Bolivar, F., Rodriguez, R. L., Greene, P. J., Betlach, M. C., Heyneker, H. L., Boyer, H. W., Crosa, J. H., and Falkow, S. (1977) Construction and characterization of new cloning vehicles. II. A multipurpose cloning system, *Gene*, **2**, 95-113.